

See discussions, stats, and author profiles for this publication at: <https://www.researchgate.net/publication/31981000>

# Acceleration of global warming due to carbon-cycle feedbacks in a coupled model

Article in *Nature* · December 2000

DOI: 10.1038/35047138 · Source: OAI

CITATIONS

2,681

READS

1,797

5 authors, including:



[Peter M. Cox](#)

University of Exeter

257 PUBLICATIONS 37,810 CITATIONS

[SEE PROFILE](#)



[Chris Jones](#)

Met Office

221 PUBLICATIONS 29,701 CITATIONS

[SEE PROFILE](#)

Some of the authors of this publication are also working on these related projects:



NERC 1.5K [View project](#)



JULES Physiology [View project](#)

## Acceleration of global warming due to carbon-cycle feedbacks in a coupled climate model

Peter M. Cox\*, Richard A. Betts\*, Chris D. Jones\*, Steven A. Spall\* & Ian J. Totterdell†

\* Hadley Centre, The Met Office, Bracknell, Berkshire RG12 2SY, UK

† Southampton Oceanography Centre, European Way, Southampton SO14 3ZH, UK

The continued increase in the atmospheric concentration of carbon dioxide due to anthropogenic emissions is predicted to lead to significant changes in climate<sup>1</sup>. About half of the current emissions are being absorbed by the ocean and by land ecosystems<sup>2</sup>, but this absorption is sensitive to climate<sup>3,4</sup> as well as to atmospheric carbon dioxide concentrations<sup>5</sup>, creating a feedback loop. General circulation models have generally excluded the feedback between climate and the biosphere, using static vegetation distributions and CO<sub>2</sub> concentrations from simple carbon-cycle models that do not include climate change<sup>6</sup>. Here we present results from a fully coupled, three-dimensional carbon-climate model, indicating that carbon-cycle feedbacks could significantly accelerate climate change over the twenty-first century. We find that under a 'business as usual' scenario, the terrestrial biosphere acts as an overall carbon sink until about 2050, but turns into a source thereafter. By 2100, the ocean uptake rate of 5 Gt C yr<sup>-1</sup> is balanced by the terrestrial carbon source, and atmospheric CO<sub>2</sub> concentrations are 250 p.p.m.v. higher in our fully coupled simulation than in uncoupled carbon models<sup>2</sup>, resulting in a global-mean warming of 5.5 K, as compared to 4 K without the carbon-cycle feedback.

The general circulation model (GCM) that we used is based on the third Hadley Centre coupled ocean-atmosphere model, HadCM3<sup>7</sup>, which we have coupled to an ocean carbon-cycle model (HadOCC) and a dynamic global vegetation model (TRIFFID). The atmospheric physics and dynamics of our GCM are identical to those used in HadCM3, but the additional computational expense of including an interactive carbon cycle made it necessary to reduce the ocean resolution to 2.5° × 3.75°, necessitating the use of flux adjustments in the ocean component to counteract climate drift. HadOCC accounts for the atmosphere-ocean exchange of CO<sub>2</sub>, and the transfer of CO<sub>2</sub> to depth through both the solubility pump and the biological pump<sup>8</sup>. TRIFFID models the state of the biosphere in terms of the soil carbon, and the structure and coverage of five functional types of plant within each model gridbox (broadleaf tree, needleleaf tree, C<sub>3</sub> grass, C<sub>4</sub> grass and shrub). Further details on HadOCC and TRIFFID are given in Methods.


The coupled climate/carbon-cycle model was brought to equilibrium with a 'pre-industrial' atmospheric CO<sub>2</sub> concentration of 290 p.p.m.v., starting from an observed landcover data set<sup>9</sup>. The resulting state was stable, with negligible net land-atmosphere and



ocean–atmosphere carbon fluxes in the long-term mean, and no discernible drift in atmospheric  $\text{CO}_2$  concentration. This simulation produces the locations of the main land biomes, and estimates of ocean carbon (38,100 Gt C), vegetation carbon (493 Gt C), soil carbon (1,180 Gt C) and terrestrial net primary productivity (60 Gt  $\text{C yr}^{-1}$ ) that are within the range of other estimates<sup>2,10–12</sup>. Ocean primary productivity is also compatible with results derived from remote sensing<sup>13,14</sup>, producing a global-mean total of 53 Gt  $\text{C yr}^{-1}$ , and realistic seasonal and latitudinal variations<sup>15</sup>.

The simulated carbon cycle displays significant interannual variability, which is driven by the model-generated El Niño/Southern Oscillation (ENSO). A realistic response to internal climate variability is an important prerequisite for any carbon-cycle model to be used in climate change predictions. Fluctuations in annual-mean atmospheric  $\text{CO}_2$  are correlated with the phase of ENSO, as indicated by the Nino3 index (Fig. 1). During El Niño conditions (positive Nino3), the model simulates an increase in atmospheric  $\text{CO}_2$ ; this increase results from the terrestrial biosphere acting as a large source (especially in Amazonia<sup>16</sup>), which is only partially offset by a reduced outgassing from the tropical Pacific Ocean. The opposite is true during the La Niña phase. The overall sensitivity of the modelled carbon cycle to ENSO variability is consistent with the observational record<sup>17</sup>, demonstrating that the coupled system responds realistically to climate anomalies.


Transient simulations were carried out for 1860–2100, using  $\text{CO}_2$  emissions as given by the IS92a scenario<sup>18</sup>. Other greenhouse gases were also prescribed from IS92a, but the radiative effects of sulphate aerosols were omitted. Three separate runs were completed to isolate the effect of climate/carbon-cycle feedbacks; an experiment with prescribed IS92a  $\text{CO}_2$  and fixed vegetation (that is, a ‘standard’ GCM climate change simulation), an experiment with interactive  $\text{CO}_2$  and dynamic vegetation but no direct effects of  $\text{CO}_2$  on climate (akin to ‘offline’ carbon-cycle projections that neglect climate change<sup>6</sup>), and a fully coupled climate/carbon-cycle simulation.



**Figure 1** Modelled and observed interannual variability in the atmospheric  $\text{CO}_2$  concentration. The figure shows the anomaly in the growth rate of atmospheric  $\text{CO}_2$  versus the Nino3 index, taken from our pre-industrial control simulation (crosses) and the Mauna Loa observations (triangles). (The Nino3 index is the annual mean sea surface temperature anomaly in the tropical Pacific, 150° W–90° W, 5° S–5° N.) The gradients of the dashed and dot-dashed lines represent the sensitivity of the carbon cycle to ENSO, as given by the observations and the model, respectively. We have excluded observations that immediately follow major volcanic events (data points shown by diamonds), since during these years the carbon cycle may have been significantly perturbed by the induced tropospheric cooling.

Figure 2 shows results from the fully coupled run. From 1860 to 2000, the simulated stores of carbon in the ocean and on land increase by about 100 Gt C and 75 Gt C, respectively. However, the atmospheric  $\text{CO}_2$  is 15–20 p.p.m.v. too high by the present day (corresponding to a timing error of about 10 years). Possible reasons for this include an overestimate of the prescribed net land-use emissions and the absence of other important climate forcing factors. The modelled global mean temperature increase from 1860 to 2000 is about 1.4 K (Fig. 3b), which is higher than observed<sup>19</sup>, probably due to the absence of cooling from anthropogenic aerosols<sup>20</sup>. Offline tests suggest that such a relative warming can suppress the terrestrial carbon sink by enhancing soil and plant respiration<sup>11</sup>. Nevertheless, the rate of increase of  $\text{CO}_2$  from 1950 to 2000 closely follows the recent observational record, which implies that the airborne fraction is being well simulated over this period. For the 20 years centred on 1985, the mean land and ocean uptake of carbon are 1.5 and 1.6 Gt  $\text{yr}^{-1}$ , respectively (compare best estimates for the 1980s of  $1.8 \pm 1.8$  and  $2.0 \pm 0.8$  Gt  $\text{yr}^{-1}$ )<sup>2</sup>. The model therefore captures the most important characteristics of the present-day carbon cycle.

The simulated atmospheric  $\text{CO}_2$  diverges much more rapidly from the standard IS92a concentration scenario in the future. First, vegetation carbon in South America begins to decline, as a drying and warming of Amazonia initiates loss of forest (Fig. 4a). This is driven purely by climate change, as can be seen by comparing the fully coupled run (red lines) to the run without global warming (blue lines). The effects of anthropogenic deforestation on land-cover are neglected in both cases. A second critical point is reached at about 2050, when the land biosphere as a whole switches from being a weak sink for  $\text{CO}_2$  to being a strong source (Fig. 2). The reduction in terrestrial carbon from around 2050 onward is associated with a widespread climate-driven loss of soil carbon (Fig. 4b). An increase in the concentration of atmospheric  $\text{CO}_2$  alone tends to increase the rate of photosynthesis and thus terrestrial carbon storage, provided that other resources are not limiting<sup>4</sup>. However, plant maintenance and soil respiration rates both increase with temperature. As a consequence, climate warming (the indirect effect of a  $\text{CO}_2$  increase) tends to reduce terrestrial carbon storage<sup>11</sup>, especially in the warmer regions where an increase in temperature is



**Figure 2** Budgets of carbon during the fully coupled simulation. The thick line shows the simulated change in atmospheric  $\text{CO}_2$ . The thinner lines show the integrated impact of the emissions, and of land and ocean fluxes, on the atmospheric  $\text{CO}_2$  increase, with negative values implying net uptake of  $\text{CO}_2$ . We note that the terrestrial biosphere takes up  $\text{CO}_2$  at a decreasing rate from about 2010 onwards, becoming a net source at around 2050. By 2100 this source from the land almost balances the oceanic sink, so that atmospheric carbon content is increasing at about the same rate as the integrated emissions (that is, the airborne fraction is  $\sim 1$ ).




not beneficial for photosynthesis. At low  $\text{CO}_2$  concentrations the direct effect of  $\text{CO}_2$  dominates, and both vegetation and soil carbon increase with atmospheric  $\text{CO}_2$ . But as  $\text{CO}_2$  increases further, terrestrial carbon begins to decrease, because the direct effect of  $\text{CO}_2$  on photosynthesis saturates but the specific soil respiration rate continues to increase with temperature. The transition between these two regimes occurs abruptly at around 2050 in this experiment (Fig. 4b). The carbon stored on land decreases by about 170 Gt C from 2000 to 2100, accelerating the rate of atmospheric  $\text{CO}_2$  increase over this period.

The ocean takes up about 400 Gt C over the same period, but at a rate which is asymptotically approaching  $5 \text{ Gt C yr}^{-1}$  by 2100. This reduced efficiency of oceanic uptake is partly a consequence of the nonlinear dependence of the partial pressure of dissolved  $\text{CO}_2$  on the total ocean carbon concentration, but may also have contributions from climate change<sup>3</sup>. Although the thermohaline circulation of the ocean weakens<sup>21</sup> by about 25% from 2000 to 2100, this is much less of a reduction than seen in some previous simulations<sup>22</sup>, and the corresponding effect on ocean carbon uptake is less significant. In our experiment, increased thermal stratification due to warming of the sea surface suppresses upwelling, which reduces nutrient availability and lowers primary production by about 5%. However, ocean-only tests suggest a small effect of climate change on oceanic carbon uptake, as this reduction in the

biological pump is compensated by reduced upwelling of deep waters which have high concentrations of total carbon.


By 2100 the modelled  $\text{CO}_2$  concentration is about 980 p.p.m.v. in the coupled experiment, which is more than 250 p.p.m.v. higher than the standard IS92a scenario or that simulated in the 'offline' experiment (Fig. 3a). As a result, the global-mean land temperatures increase from 1860 to 2100 by about 8 K, rather than the 5.5 K of the standard scenario (Fig. 3b).

These numerical experiments demonstrate the potential importance of climate/carbon-cycle feedbacks, but the magnitude of these in the real Earth system is still highly uncertain. Terrestrial carbon models differ in their responses to climate change<sup>11,12</sup>, owing to gaps in basic understanding of processes. For example, the potential conversion of the global terrestrial carbon sink to a source is critically dependent upon the long-term sensitivity of soil respiration to global warming, which is still a subject of debate<sup>23</sup>. The experiments presented here exclude the potentially large direct human influences on terrestrial carbon uptake through changes in landcover and land management. Local effects, such as the possible climate-driven loss of the Amazon rainforest, rest upon uncertain aspects of regional climate change, and may be 'short-circuited' by direct human deforestation. A full assessment of the uncertainties must await further coupled experiments utilizing alternative representations of processes and including a more complete set of natural



**Figure 3** Effect of climate/carbon-cycle feedbacks on  $\text{CO}_2$  increase and global warming. **a**, Global-mean  $\text{CO}_2$  concentration, and **b**, global-mean and land-mean temperature, versus year. Three simulations are shown; the fully coupled simulation with interactive  $\text{CO}_2$  and dynamic vegetation (red lines), a standard GCM climate change simulation with

prescribed (IS92a)  $\text{CO}_2$  concentration and fixed vegetation (dot-dashed lines), and the simulation which neglects direct  $\text{CO}_2$ -induced climate change (blue lines). The slight warming in the latter is due to  $\text{CO}_2$ -induced changes in stomatal conductance and vegetation distribution.



**Figure 4** Effect of global warming on changes in land carbon storage. The red lines represent the fully coupled climate/carbon-cycle simulation, and the blue lines are from the 'offline' simulation which neglects direct  $\text{CO}_2$ -induced climate change. The figure

shows simulated changes in vegetation carbon (**a**) and soil carbon (**b**) for the global land area (continuous lines) and South America alone (dashed lines).



and anthropogenic forcing factors (for example, land-use change, forest fires, sulphate aerosol concentrations and nitrogen deposition). However, our results indicate that it will be essential to accurately represent previously neglected carbon-cycle feedbacks if we are to successfully predict climate change over the next 100 years. □

## Methods

### Ocean carbon-cycle model

The inorganic component of HadOCC has been extensively tested as part of the Ocean Carbon Cycle Intercomparison Project; it was found to reproduce tracer distributions to an accuracy consistent with other ocean GCMs<sup>24</sup>. The biological component treats four additional ocean fields: nutrient, phytoplankton, zooplankton and detritus<sup>8</sup>. The phytoplankton population changes as a result of the balance between growth, which is controlled by light level and the local concentration of nutrient, and mortality, which is mostly as a result of grazing by zooplankton. Detritus, which is formed by zooplankton excretion and by phyto- and zooplankton mortality, sinks at a fixed rate and slowly remineralizes to reform nutrient and dissolved inorganic carbon. Thus both nutrient and carbon are absorbed by phytoplankton near the ocean surface, pass up the food chain to zooplankton, and are eventually remineralized from detritus in the deeper ocean. The model also includes the formation of calcium carbonate and its dissolution at depth (below the lysocline).

### Terrestrial carbon-cycle model

TRIFFID (top-down representation of interactive foliage and flora including dynamics) has been used offline in a comparison of dynamic global vegetation models<sup>11</sup>. Carbon fluxes for each vegetation type are calculated every 30 minutes as a function of climate and atmospheric CO<sub>2</sub> concentration, from a coupled photosynthesis/stomatal-conductance scheme<sup>25,26</sup>, which utilizes existing models of leaf-level photosynthesis in C<sub>3</sub> and C<sub>4</sub> plants<sup>27,28</sup>. The accumulated fluxes are used to update the vegetation and soil carbon every 10 days. The natural landcover evolves dynamically based on competition between the vegetation types, which is modelled using a Lotka-Volterra approach and a tree-shrub-grass dominance hierarchy. We also prescribe some agricultural regions; in which grasslands are assumed to be dominant. Carbon lost from the vegetation as a result of local litterfall or large-scale disturbance is transferred into a soil carbon pool, where it is broken down by microorganisms that return CO<sub>2</sub> to the atmosphere. The soil respiration rate is assumed to double for every 10 K of warming<sup>29</sup>, and is also dependent on the soil moisture content<sup>30</sup>. Changes in the biophysical properties of the land surface<sup>5</sup>, as well as changes in terrestrial carbon, feed back onto the atmosphere.

Received 6 January; accepted 26 September 2000.

- Houghton, J. T. et al. (eds) *Climate Change 1995: The Science of Climate Change* (Cambridge Univ. Press, Cambridge, 1996).
- Schimel, D. et al. in *Climate Change 1995: The Science of Climate Change* Ch. 2 (eds Houghton, J. T. et al.) 65–131 (Cambridge Univ. Press, Cambridge, 1995).
- Sarmiento, J., Hughes, T., Stouffer, R. & Manabe, S. Simulated response of the ocean carbon cycle to anthropogenic climate warming. *Nature* **393**, 245–249 (1998).
- Cao, M. & Woodward, F. I. Dynamic responses of terrestrial ecosystem carbon cycling to global climate change. *Nature* **393**, 249–252 (1998).
- Betts, R. A., Cox, P. M., Lee, S. E. & Woodward, F. I. Contrasting physiological and structural vegetation feedbacks in climate change simulations. *Nature* **387**, 796–799 (1997).
- Enting, I., Wigley, T. & Heimann, M. *Future Emissions and Concentrations of Carbon Dioxide; Key Ocean/Atmosphere/Land Analyses* (Technical Paper 31, Division of Atmospheric Research, CSIRO, Melbourne, 1994).
- Gordon, C. et al. The simulation of SST, sea ice extents and ocean heat transports in a version of the Hadley Centre coupled model without flux adjustments. *Clim. Dyn.* **16**, 147–168 (2000).
- Palmer, J. R. & Totterdell, I. J. Production and export in a global ocean ecosystem model. *Deep-Sea Res.* (in the press).
- Wilson, M. F. & Henderson-Sellers, A. A global archive of land cover and soils data for use in general circulation climate models. *J. Clim.* **5**, 119–143 (1985).
- Zinke, P. J., Stangenberger, A. G., Post, W. M., Emanuel, W. R. & Olson, J. S. *Worldwide Organic Soil Carbon and Nitrogen Data* (NDP-018, Carbon Dioxide Information Center, Oak Ridge National Laboratory, Oak Ridge, Tennessee, 1986).
- Cramer, W. et al. Global response of terrestrial ecosystem structure and function to CO<sub>2</sub> and climate change: results from six dynamic global vegetation models. *Glob. Change Biol.* (in the press).
- VEMAP Members. Vegetation/ecosystem modelling and analysis project: comparing biogeography and biogeochemistry models in a continental-scale study of terrestrial responses to climate change and CO<sub>2</sub> doubling. *Glob. Biogeochem. Cycles* **9**, 407–437 (1995).
- Longhurst, A., Sathyendranath, S., Platt, T. & Caverhill, C. An estimate of global primary production in the ocean from satellite radiometer data. *J. Plank. Res.* **17**, 1245–1271 (1995).
- Field, C., Behrenfeld, M., Randerson, J. & Falkowski, P. Primary production of the biosphere: integrating terrestrial and oceanic components. *Science* **281**, 237–240 (1998).
- Antoine, D., Andre, J.-M. & Morel, A. Oceanic primary production 2. Estimation at global scale from satellite (Coastal Zone Color Scanner) chlorophyll. *Glob. Biogeochem. Cycles* **10**, 57–69 (1996).
- Tian, H. et al. Effects of interannual climate variability on carbon storage in Amazonian ecosystems. *Nature* **396**, 664–667 (1998).
- Keeling, C. D., Whorf, T., Whalen, M. & der Plicht, J. V. Interannual extremes in the rate of rise of atmospheric carbon dioxide since 1980. *Nature* **375**, 666–670 (1995).
- Houghton, J. T., Callander, B. A. & Varney, S. K. (eds) *Climate Change 1992: The Supplementary Report to the IPCC Scientific Assessment* (Cambridge Univ. Press, Cambridge, 1992).

- Nicholls, N. et al. in *Climate Change 1995: The Science of Climate Change* Ch. 3 (eds Houghton, J. T. et al.) (Cambridge Univ. Press, Cambridge, 1996).
- Mitchell, J. F. B., Johns, T. C., Gregory, J. M. & Tett, S. F. B. Climate response to increasing levels of greenhouse gases and sulphate aerosols. *Nature* **376**, 501–504 (1995).
- Wood, R. A., Keen, A. B., Mitchell, J. F. B. & Gregory, J. M. Changing spatial structure of the thermohaline circulation in response to atmospheric CO<sub>2</sub> forcing in a climate model. *Nature* **399**, 572–575 (1999).
- Sarmiento, J. & Quere, C. L. Oceanic carbon dioxide uptake in a model of century-scale global warming. *Nature* **274**, 1346–1350 (1996).
- Giardina, C. & Ryan, M. Evidence that decomposition rates of organic carbon in mineral soil do not vary with temperature. *Nature* **404**, 858–861 (2000).
- Orr, J. C. in *Ocean Storage of Carbon Dioxide, Workshop 3: International Links and Concerns* (ed. Ormerod, W.) 33–52 (IEA R&D Programme, CRE Group Ltd, Cheltenham, UK, 1996).
- Cox, P. M., Huntingford, C. & Harding, R. J. A canopy conductance and photosynthesis model for use in a GCM land surface scheme. *J. Hydrol.* **212**, 79–94 (1998).
- Cox, P. M. et al. The impact of new land surface physics on the GCM simulation of climate and climate sensitivity. *Clim. Dyn.* **15**, 183–203 (1999).
- Collatz, G. J., Ball, J. T., Grievet, C. & Berry, J. A. Physiological and environmental regulation of stomatal conductance, photosynthesis and transpiration: a model that includes a laminar boundary layer. *Agric. Forest Meteorol.* **54**, 107–136 (1991).
- Collatz, G. J., Ribas-Carbo, M. & Berry, J. A. A coupled photosynthesis-stomatal conductance model for leaves of C<sub>4</sub> plants. *Aust. J. Plant Physiol.* **19**, 519–538 (1992).
- Raich, J. & Schlesinger, W. The global carbon dioxide flux in soil respiration and its relationship to vegetation and climate. *Tellus B* **44**, 81–99 (1992).
- McGuire, A. et al. Interactions between carbon and nitrogen dynamics in estimating net primary productivity for potential vegetation in North America. *Glob. Biogeochem. Cycles* **6**, 101–124 (1992).

## Acknowledgements

We thank J. Mitchell and G. Jenkins for comments on earlier versions of the manuscript. This work was supported by the UK Department of the Environment, Transport and the Regions.

Correspondence and requests for materials should be addressed to P.M.C. (e-mail: pmcox@meto.gov.uk).